

Thermodynamic witness of quantum probing

H.Dong,¹ X.F. Liu,² and C.P. Sun^{1,*}

¹*Institute of Theoretical Physics, Chinese Academy of Science, Beijing 100080, China*

²*Department of Mathematics, Peking University, Beijing 100871, China*

The thermodynamic influence of quantum probing on an object is studied. Here, quantum probing is understood as a pre-measurement based on a non-demolition interaction, which records some information of the probed object, but does not change its energy state when both the probing apparatus and the probed object are isolated from the environment. It is argued that when the probing apparatus and the probed object are immersed in a same equilibrium environment, the probing can affect the effective temperature of the object or induce a quantum isothermal process for the object to transfer its energy. This thermodynamic feature can be regarded as a witness of the existence of quantum probing even if the quantum probing would not disturb the object if the environment were not present.

PACS numbers: 03.65.Ta, 03.65.Yz, 05.30.Ch

Introduction- The Landauer's principle that the erasure of one bit of information requires a minimum heat generation of $k_B T \ln 2$, which is based on the second law of thermodynamics [1, 2], underlies the thermodynamics of information processing. This principle eventually resolves the Maxwell's demon paradox: why a demon can assist a binary thermal medium to do extra work [2]. For the science and technology of quantum information, the Landauer's principle is undoubtedly crucial since it gives a physical limitation on the spatial-time scales of logical devices on chips.

However, in the arguments *for* and *against* the Landauer's principle, the conventional question in the thermodynamics of information processing that whether the measurement process requires a cost of heat generation has never been convincingly answered. Surely, earlier authors had touched on this problem, but it has not been clarified yet because the measurement process is not defined properly. Particularly, in the quantum approach of measurement, people can not be unanimous for some fundamental problems, such as whether or not there exist wave function collapse [3]. In fact, to clarify the situation, we need to answer the following subtle questions in an unambiguous way: when can we say a system is performing a measurement on another system and what kind of measurement can dissipate information?

In this paper, we refer quantum probing to a pre-measurement [4, 5] based on the non-demolition coupling of the probed system S to the apparatus A , which is only a unitary process to produce entanglement between S and A , and does not concern the subtle, seemingly philosophical arguments, such as wave function collapses. Generally speaking, when the energy state of the measured system S is not influenced by the coupling to the measuring apparatus A , but A can record some information of the system S , the measurement performed

by A on S is called a pre-measurement. The so called non-demolition (pre-) measurement [4] is an ideal measurement under some circumstances. In the case of non-demolition measurement, the system-apparatus coupling V_{SA} commutes with the Hamiltonian H_S of the measured system, but does not commute with the Hamiltonian H_A of the measuring apparatus. When studying non-demolition measurement it is usually assumed that both A and S are isolated from an environment. In this paper, we will investigate the effect of environment on the measured system in a non-demolition measurement. In the following discussion, we will use the name quantum probing or just probing when regarding such non-demolition measurement. In the presence of environment, the practical thermalization [6] of the total system $A + S$ will have thermodynamic effects on S . These effects can be regarded as the thermodynamic witness of quantum probing.

When the existence of an environment E is considered, quantum probing is understood in two steps (illustrated in Fig. 1) :

- Due to the extreme-weakness of the coupling between $S + A$ and E , it can be neglected within the dephasing time τ_2 of the system, the dephasing being the result of the interaction between S and A . In the end of this step, the initial factorized state of $S + A$ becomes a state assuming the form of an ideal Schmidt decomposition $\sum c_n |n\rangle \otimes |D_n\rangle$ [5], with $\langle D_m | D_n \rangle = \delta_{mn}$.
- If the probing apparatus continues to probe the state of the system S , the non-demolition coupling should hold for a longer time, and the communication with the environment will result in the thermalization of the total system $S + A$. In this step, the information of the initial state should be erased totally, but the correlation between A and S needs to exist to leave the witness of quantum probing.

As shown as follows, it is the common environment of the probed system and the probing apparatus that

*Electronic address: suncp@itp.ac.cn;
URL: <http://power.itp.ac.cn/~suncp/>




FIG. 1: Time scale of the non-demolition pre-measurement and thermalization. (1) In the time interval $(0, \tau_2)$, the apparatus makes the pre-measurement: $\sum c_n |n\rangle \otimes |D\rangle \rightarrow \sum c_n |n\rangle \otimes |D_n\rangle$ and the environment does not play the role, since its coupling to $S + A$ is weaker than that between S and A . (2) In the time interval (τ_2, τ_1) , the effect of thermalization due to the environment becomes prominent. The total system $S + A$ is finally thermalized in the canonical state $\rho_{can}(T)$ with temperature T .

selects a special set of ideal entanglement states of $S + A$ to be thermalized, so that these witnesses of quantum probing are of thermodynamics, and thus observable in the classical or macroscopic level.

Universal Setup for quantum probing apparatus- Let A be a general apparatus weakly coupling to the system S to be probed. We require that the energy spectrum of A be denser than that of S . The following heuristic argument may help to justify this requirement: to measure the spatial scale of an object, the ruler should have a much finer graduation than the size of this object.

To be precise, let $H_A = \sum_k \epsilon_k |k\rangle \langle k|$ be the spectrum decomposition of the Hamiltonian of A and $H_S = \sum_n E_n |n\rangle \langle n|$ the spectrum decomposition of the Hamiltonian of S , the requirement then can be expressed as $\min\{|E_n - E_{n+M}|\} \gg \max\{|\epsilon_k - \epsilon_{k+1}|\}$. Here, $|k\rangle$ is the eigenvector of H_A corresponding to the eigenvalue ϵ_k , and $|n\rangle$ is the eigenvector of H_S corresponding to the eigenvalue E_n . Let V_{AS} be a weak coupling between S and A . The weakness of V_{AS} means that its effect on the dynamics of the total system $S + A$ can be well studied by the perturbation method [7].

To investigate the behavior of the total system $S + A$ immersed in an environment, we consider the partition function

$$Z = \text{Tr}(e^{-\beta H}) = \text{Tr}(W^\dagger e^{-\beta H} W)$$

Here, as a trick, we have introduced a unitary transformation $W = \exp(-S)$, defined by an anti-Hermitian operator S , which is a perturbation quantity of the same order as V_{AS} . If

$$V_{AS} + [H_A + H_S, S] = 0,$$

then the partition function can be approximated as $Z = \text{Tr} \exp(-\beta H_{\text{eff}})$ where the effective Hamiltonian

$$H_{\text{eff}} = H_A + H_S + \frac{1}{2} [V_{AS}, S], \quad (1)$$

is just the Frohlich-Nakajima Hamiltonian in solid state physics [7]. Here, we re-derive it to justify its applicability in thermodynamics.

Since the minimal energy level spacing of the system is much larger than the energy level spacing of the apparatus, the effective interaction V_{eff} can be obtained as $V_{\text{eff}} = \sum_n \mathcal{H}(n) |n\rangle \langle n|$ where

$$\mathcal{H}(n) = \sum_k G(n) |k\rangle \langle k'| \quad (2)$$

is a branched effective Hamiltonian of the apparatus corresponding to the situation that the system is prepared in the state $|n\rangle$ with the coupling

$$G(n) = \frac{\langle nk | V_{AS} | nk' \rangle}{(\epsilon_k - \epsilon_{k'})}. \quad (3)$$

The above obtained effective interaction V_{eff} satisfies $[H_S, V_{\text{eff}}] = 0$. Thus the total effective Hamiltonian $H_{\text{eff}} = H_S + H_A + V_{\text{eff}}$ describes a non-demolition measurement without the presence of an environment. In this case, the factorized initial state $|\varphi(0)\rangle = \sum_n c_n |n\rangle \otimes |D\rangle$ will evolve into $|\varphi(t)\rangle = \sum_n c_n |n\rangle \otimes |D_n(t)\rangle$ where $|D_n(t)\rangle$ takes the form $|D_n(t)\rangle = \exp[-iG(n)t] |D\rangle$. If $\{|D_n(t)\rangle\}$ becomes an orthogonal set when t approaches infinity, then the time evolution $|\varphi(t)\rangle$ represents a process of ideal pre-measurement.

Thermodynamic effects of measurements- Now we study the change in the thermodynamic features of the system caused by the apparatus. We assume that both the probed system and the measuring apparatus are immersed in the same thermal bath with temperature T or inverse temperature $\beta = 1/k_B T$. After or during the measuring process, the total system $S + A$ will reach the state with the same temperature T , if the total system is non-degenerate. Then we can calculate the reduced density matrix $\rho_S = \text{Tr}_A [\exp(-\beta(H_S + H_A + V_{\text{eff}}))]$, obtaining

$$\rho_S = \frac{1}{Z'_S} \sum_n e^{-\beta E_n} \xi(n) |n\rangle \langle n|, \quad (4)$$

where $Z'_S = \sum_n \exp(-\beta E_n) \xi(n)$ and

$$\xi(n) = \text{Tr}_A \{ \langle n | \exp[-\beta(H_A + V_{\text{eff}})] | n \rangle \} \quad (5)$$

is a formal factor depending on the system state $|n\rangle$ and vanishing trivially when no coupling exists.

Here, we consider a manipulation process. (i) Initially, no probing apparatus is coupled to the system, which is in equilibrium with the heat bath with temperature T . (ii) Then, the apparatus begin to probe the system at time $t = 0$. As the evolution described in Fig. 2, the total system $S + A$ reaches the state with the same temperature T . We observe that the thermodynamic effect of measurement implied in Eq. (6) allows two interpretations. These two interpretations are illustrated in Fig. 2 as those in Ref. [8, 9].

The first interpretation goes as follows. The change from the close thermal state of the system $\rho_{Sc} = \exp(-\beta H_S) / Z_S$ to the modified thermal state ρ_S is understood as a quantum isometric process, represented by




FIG. 2: Two interpretations. (i) Isometric process ($A \rightarrow C$, the solid line): the inverse temperature β is fixed in the process. (ii) Isothermal process ($A \rightarrow B$, the dotted line): The level spacing Δ is fixed in the process.

the solid line between the points A and C in Fig. 2. In this process, the temperature keeps unchanged, but the energy level spacings are alternated by the interaction with the apparatus. Accordingly, we define $\xi(n) = \exp(-\beta \Delta E_n)$, then ρ_S can be written in the form with clear physical meaning:

$$\rho_S = \frac{1}{Z'_S} \sum_n e^{-\beta(E_n + \Delta E_n)} |n\rangle \langle n|. \quad (6)$$

In the present case, it is rather natural to regard the inner energy change $\Delta U = \text{Tr}_S [H_S (\rho_S - \rho_{Sc})]$ as a witness of the thermodynamic role of measurement.

In the second interpretation the change from ρ_{Sc} to ρ_S is understood as a quantum isothermal process, represented by the dotted line between the points A and B in Fig. 2. In this process, the energy level spacings are fixed. To justify our using the term “isothermal process” here, at least to some extent, we define the effective temperature [6] via $\beta(n) = \ln(P_n/P_{n+1})/\Delta_n$ or

$$\beta(n) = \beta + \frac{1}{\Delta_n} \ln \frac{\xi(n)}{\xi(n+1)}, \quad (7)$$

where $\Delta_n = E_{n+1} - E_n$ is the n -th energy level spacing. We notice that generally this generalized temperature cannot be regarded as an effective temperature since it depends on the energy levels of the system. But if $\beta(n+1) - \beta(n) = 0$, i.e.,

$$\left[\frac{\xi(n+1)}{\xi(n+2)} \right]^{\Delta_n} = \left[\frac{\xi(n)}{\xi(n+1)} \right]^{\Delta_{n+1}}, \quad (8)$$

then $\beta(n)$ becomes a well defined thermodynamic parameter independent of n , which we denote by β_{eff} . In this case, the above defined generalized temperature allows the physical interpretation of effective temperature, and we have $\rho_S = \sum_n \exp(-\beta_{\text{eff}} E_n) |n\rangle \langle n| / Z'_S$, and the inner energy change

$$\Delta U = \sum_n E_n [e^{-\beta_{\text{eff}} E_n} - e^{-\beta E_n}] \quad (9)$$

reflects the thermalization effect.

The quantum role of measurement and renormalization- As an explicit illustration, we model the apparatus with weak couplings to the probed system as a collection of harmonic oscillators. According to the results of Ref. [10] and Ref. [11] (where the arguments are carried out for the bath modeling, but can work well for our setup for the apparatus), the coupling of the system to the apparatus is linear with respect to the coordinates of the bath harmonic oscillators. Let $b_j^\dagger$ (b_j) be the creation(annihilation) operator of the j -th mode of the bath with eigen-frequency ω_j , and $\lambda_n g_j$ the coupling coefficient of the system state $|n\rangle$ to the j -th mode. Then, the total Hamiltonian is obtained as

$$H = \sum_n E_n |n\rangle \langle n| + \sum_k \omega_k b_k^\dagger b_k + \sum_n \lambda_n |n\rangle \langle n| \sum_k (g_k a_k^\dagger + \text{h.c.}). \quad (10)$$

In this case, the above defined generalized inverse temperature reads

$$\beta(n) = \beta \left(1 - \frac{|\lambda_{n+1}|^2 - |\lambda_n|^2}{E_{n+1} - E_n} \varepsilon \right), \quad (11)$$

where

$$\varepsilon = \sum_k \frac{|g_k|^2}{\omega_k} = \int \rho(\omega) \frac{|g_k|^2}{\omega_k} dk \quad (12)$$

represents the self-energy of the apparatus, which causes the Lamb shift of its coupled system. As pointed above, to define reasonably an effective temperature for the system, the generalized inverse temperature $\beta(n)$ should be independent of the energy level. Here is a simple example satisfying this condition: the system is a harmonic oscillator with the energy level $E_n = (n + 1/2)\omega$ and the coupling strength $\lambda_n = \sqrt{n}$. In this example, the well defined effective inverse temperature is

$$\beta_{\text{eff}} = \beta \left(1 - \sum_k \frac{|g_k|^2}{\omega \omega_k} \right), \quad (13)$$

and the corresponding effective temperature $T_{\text{eff}} = (k_B \beta_{\text{eff}})^{-1}$ of the system is higher than that of the environment $T = (k_B \beta)^{-1}$.

For a two-level system with the excited state $|e\rangle$ and the ground state $|g\rangle$ and the energy level spacing Δ , the effective inverse temperature is also well defined. It reads

$$\beta_{\text{eff}} = \beta + \frac{1}{\Delta} \ln \frac{\xi_g}{\xi_e}. \quad (14)$$

If the apparatus is a single mode cavity with frequency ω_b and the two-level system is coupled to it by the dipole interaction with coupling strength g , then in the large detuning case $\omega_b \gg \Delta$, the formal factor ξ_e and ξ_g can be explicitly calculated as follows:

$$\xi_e = \sum_n \exp \left\{ -\beta \left[\omega_b n + \frac{|g|^2}{\omega_b - \Delta} (n+1) \right] \right\},$$




FIG. 3: Fidelity. (a) Fidelity $\mathcal{F}$ vs the level shift λ . (b) Fidelity $\mathcal{F}$ vs the temperature shift ΔT . Here, we take $\omega = 1$ and $\beta = 1$.

$$\xi_g = \sum_n \exp \left\{ -\beta \left[\omega_b n + \frac{|g|^2}{\omega_b - \Delta} n \right] \right\}. \quad (15)$$

Since $\xi_g > \xi_e$, we are led to the conclusion that the measurement will decrease the temperature of the system by

$$\Delta T = \frac{\frac{1}{\Delta} \ln \frac{\xi_g}{\xi_e}}{\beta + \frac{1}{\Delta} \ln \frac{\xi_g}{\xi_e}} T. \quad (16)$$

Finally, to quantitatively evaluate the thermalized state resulting from the interaction with the environment, let us check the fidelity [12] $\mathcal{F}$ of such a state to the ordinary canonical state $\rho_{\text{can}} = \exp(-\beta H_S) / \text{Tr}[\exp(-\beta H_S)]$. Generally for the reduced density matrix in Eq. 4, the fidelity between the initial state and the final state reads

$$\mathcal{F} = \frac{\sum_n e^{-\beta E_n} \sqrt{\xi(n)}}{(\sum_n e^{-\beta E_n})^{1/2} (\sum_n e^{-\beta E_n} \xi(n))^{1/2}}. \quad (17)$$

When the probed system is also a harmonic oscillator discussed above, the fidelity can be analytically obtained as

$$\mathcal{F} = \frac{\sqrt{\sinh \frac{\beta \omega}{2} \sinh \frac{\beta(\omega - \lambda)}{2}}}{\sinh \frac{\omega - \lambda/2}{2}}, \quad (18)$$

where $\lambda = \sum_k |g_k|^2 / \omega_k$ characterizes the shift of the energy level of the harmonic oscillator, which reflects the effect due to the coupling to the apparatus. In Fig. 3(a), the fidelity is plotted against the level shift. As the coupling between the apparatus and the system is turn on, the energy level for the system is effective $E'_n = n(\omega - \lambda)$ based on the first interpreting of the reduced density matrix. The fidelity between the canonical thermal and the reduced density matrix decreases as the coupling becomes strong. Therefore, the system gradually deviates from canonical thermal state, leaving an evidence of witness of the apparatus. For the second interpreting, we plot the fidelity $\mathcal{F}$ as the function of the temperature shift $\Delta T = 1/\beta_{\text{eff}} - 1/\beta = [\beta(\omega/\lambda - 1)]^{-1}$ in Fig. 3(b).

Conclusion- In summary, for the weak coupling case, we show from the generalized approach of Frohlich-Nakajima transformation the universality of a non-demolition Hamiltonian in connection with the probing process. Based on this general non-demolition measurement, we investigate the probing effect on the system when the total measurement happens in a reservoir. It is concluded that the probing of the system can be witted through the change of the effective temperature, even though there is no direct energy exchange between the system and the detector. To characterize the change of state for the system, we evaluate the fidelity of the modified canonical thermal state to the original canonical state without being probed.

The work is supported by National Natural Science Foundation of China and the National Fundamental Research Programs of China under Grant No. 10874091 and No. 2006CB921205.

[1] R. Landauer, IBM J. Res. Dev. **5**, 183 (1961).
 [2] H.S. Leff, & A.F. Rex(eds), *Maxwell's Demon 2: Entropy, Classical and Quantum Information, Computing*. Institute of Physics. (2003).
 [3] W. H. Zurek, Rev. Mod. Phys. **75**, 715 (2003).
 [4] V. B. Braginsky and F. Y. Khalili, *Quantum Measurement*, Cambridge University Press, (1992).
 [5] C.P. Sun, Phys. Rev. A **48**, 898 (1993).

[6] H. Dong, S. Yang, X. F. Liu, and C. P. Sun, Phys. Rev. A **76**, 044104 (2007); H. T. Quan, P. Zhang, and C. P. Sun, Phys. Rev. E **73**, 036122 (2006).
 [7] H. Fröhlich, Phys. Rev. **79**, 845(1950); S. Nakajima, Adv. Phys. **4**, 363(1953).
 [8] H. T. Quan, P. Zhang, and C. P. Sun, Phys. Rev. E **72**, 056110 (2005)
 [9] H.T. Quan, Yu-xi Liu, C. P. Sun and F. Nori, Phys.

- Rev. E **76**, 031105 (2007); H.T. Quan, Phys. Rev. E **79**, 041129 (2009)
- [10] A. O. Caldeira and A. J. Leggett, Ann. Phys. (N.Y.) **149**, 374 (1983); A. J. Leggett, S. Chakravarty, A. T. Dosey, M. P. A. Fisher, and W. Zwerger, Rev. Mod. Phys. **59**, 187 (1987).
- [11] C.P. Sun, H. Zhan and X.F. Liu, Phys.Rev. A **58**, 1810 (1998).
- [12] P. Zanardi, H. T. Quan, X. Wang, C. P. Sun Phys. Rev. A **75**, 032109 (2007) and references therein.